

After revision anterior cruciate ligament reconstruction, who returns to sport? A systematic review and meta-analysis

Alberto Grassi,¹ Stefano Zaffagnini,¹ Giulio Maria Marcheggiani Muccioli,¹ Maria Pia Neri,¹ Stefano Della Villa,² Maurilio Marcacci¹

¹Clinica Ortopedica e Traumatologica II—Lab. di Biomeccanica, ed Innovazione Tecnologica, Istituto Ortopedico Rizzoli, Bologna, Italy
²Isokinetic FIFA Medical Center of Excellence, Bologna, Italy

Correspondence to

Dr Alberto Grassi, Clinica Ortopedica e Traumatologica II—Lab. di Biomeccanica, ed Innovazione Tecnologica, Istituto Ortopedico Rizzoli, Via di Barbiano, 1/10, Bologna 40136, Italy; alberto.grassi3@studio.unibo.it

Accepted 18 May 2015

ABSTRACT

Background Return to sport and to pre-injury level represents an important outcome after both primary and revision anterior cruciate ligament (ACL) reconstructions.

Purpose The aim of the present meta-analysis was to determine the return to sport rate after revision ACL reconstruction.

Material and methods A systematic search was performed of the MEDLINE, Embase and the Cochrane Central Register of Controlled Trials Databases. All the studies that reported return to sport, return to pre-injury sport level and return to high level/competitive sport was considered for the meta-analysis. The overall pooled mean of post-operative knee laxity and pooled rate of positive pivot-shift and objective International Knee Documentation Committee (IKDC) categories was calculated as well.

Results Overall, 472 abstracts were identified and screened for inclusion and only 16 studies reported the rate of return to any level of sport activity at the final follow-up of 4.7 years (range 1.0–13.2 years), showing a pooled rate of 85.3% (CI 79.7 to 90.2). The return to pre-injury sport level was achieved in 53.4% (CI 37.8 to 68.7) of cases. Normal or quasi-normal objective IKDC, less than 5 mm of side-to-side difference at arthrometric evaluations and grade I-II pivot-shift test were reported in 84%, 88% and 93% patients, respectively.

Conclusions In spite of almost 8 patients out of 10 returning to sport after revision ACL reconstruction and showing good stability, only half of the patients returned to the same pre-injury sport level.

INTRODUCTION

Return to sport and especially to pre-injury levels represents for patients one of the most important clinical outcomes after ACL reconstruction. Despite clinical satisfaction and adequate knee range of motion (ROM), stability and muscle strength, the ability to perform sports activity is believed to represent a good indicator of overall knee health and of the ability of the joint to sustain external load and stress, particularly in high-level, competitive and pivoting activities.¹ However, in certain circumstances, the normalised objective knee status and recovered muscle strength do not correspond to the ability to perform sport, especially when psychological recovery is not reached.^{2–5}

With regard to this, the outcomes of return to sport after primary ACL reconstruction have been extensively investigated through high-level clinical studies and well-conducted meta-analyses.^{1 6 7} In contrast, the outcomes of revision ACL

reconstruction derive mostly from level IV retrospective studies.⁸ A recent meta-analysis of 21 studies investigated the main clinical outcomes of revision ACL reconstruction, reporting a return to unrestricted activities or to the previous levels of activity in 54%, but without clearly defining a precise return to sport activity definition.⁸ Moreover, since then, clinical studies reporting the results after revision ACL have almost doubled, underlining the increasing interest in this surgical procedure. If the increased number of ACL lesions, estimated at around 36.9/100 000 person-year,⁹ and the failure rate of primary ACL reconstruction of 7–10%¹⁰ are considered, it is not difficult to understand the burden of the revision ACL reconstruction procedure and its importance in the field of sports medicine and traumatology.

Nowadays ACL revision surgery, once considered but a salvage procedure, is increasingly obtaining results which are more and more similar to primary reconstruction mainly regarding clinical outcomes and knee stability,^{8 11} owing to technical surgical advances. However, the ability to perform sports activity after revision ACL reconstruction remains a not adequately investigated issue, as single studies reported the return to pre-injury sport level ranging from 0% to 100%.^{12 13}

The primary aim of the present meta-analysis was to synthesise the current evidence related to sports activity after revision ACL reconstruction, and to determine the overall average postoperative return-to-sports and return to pre-injury sport level rates. The secondary aim was to summarise key objective knee function measures, including the International Knee Documentation Committee (IKDC) score, the anteroposterior knee laxity and pivot shift test.

MATERIAL AND METHODS

Study design

A meta-analysis was performed in accordance with the PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses.¹⁴ No protocol exists for this study, which was exempt from review by Ethical Committee approval for only those studies that involved human participants, as no patients were enrolled in the study and nor were medical or surgical treatments directly administered. No external funding supported the present study.

Search strategy

An electronic search was performed on 31 September 2014 of the MEDLINE, EMBASE and

To cite: Grassi A, Zaffagnini S, Marcheggiani Muccioli GM, et al. *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2014-094089

Cochrane Central Register of Controlled Trials databases. The search terms used were “Anterior cruciate ligament” and “ACL” combined with the terms “revision”, “reoperation” and “failure”. Filters were utilised to access only those studies that investigated participants and were reported in English. No limitations were imposed on the date of publication. The reference section of each full-text obtained was scanned and hand-searched for other possible inclusions to this study, which were not identified in the electronic database search.

Eligibility criteria

Studies were included if they met the following criteria: (1) randomised controlled trial (RCT), prospective or retrospective study, (2) the study reported the outcomes of both first or multiple revision ACL reconstruction, (3) the study reported at least one of: the return to any kind of sport rate, return to pre-injury sport level. When “return to pre-injury activity” was used as the outcome without mentioning the nature of the activity, the definition was considered too vague and therefore the study was excluded. The authors of the studies that met some but not all of the criteria were contacted for further information and to clarify whether the study was eligible or not.

Study selection

Two authors (AG and GMMM) independently reviewed the studies found during the database search. Articles were screened for eligibility based on a review of the title and abstract only. Any potential disagreement was resolved via consensus discussion between two independent reviewers, with a third reviewer (SZ) being consulted if the disagreement could not be resolved. When the eligibility was unclear from the title and abstract, the full text was obtained and assessed and again evaluated for eligibility.

Data collection and analysis

All the data regarding the names of authors, year of publication, number of patients treated, final follow-up duration and graft used for revision procedure were obtained and tabulated in an Excel extraction form to systematically extract the data. Regarding the sport-specific outcomes, details of all the patients who were still practising any kind of sport at final follow-up, and of all the patients who at final follow-up returned to the same level practised before the first ACL injury, were accurately extracted from each study and evaluated as primary outcomes. When possible, the subgroups of patients who returned to high level and competitive sport activity—according to clinical scores and definitions provided in each study—were also extracted.

The secondary outcomes relative to the final follow-up were the objective IKDC^{15 16} classification, knee laxity measured as side-to-side difference at the manual maximum displacement test measured in mm with KT-1000 or KT-2000 and results of the pivot-shift test. The percentage of participants with knee classified as category A (normal knee), B (nearly normal knee), C (abnormal knee) or D (severely abnormal knee) according to the IKDC form was calculated. Arithmetic averages of knee laxity and percentages of patients with a <3, 3–5 or >5 mm side to side difference at the manual maximum displacement test were calculated as well. The total number of patients with a pivot-shift test graded as normal (equal to the contralateral side), grade I (or glide, in case defined as a small and gentle sliding reduction), grade II (or clunk, in case of reduction movement easily appreciated) and grade III (gross, in case of an abnormal movement with a pronounced clunk)^{17 18} was calculated and expressed as a percentage for each groups. As for

multiple revisions, the data regarding first revision were collected when available; otherwise, only data of the last revision were used for analysis.

Quality assessment

The studies included were assessed using a six-item checklist developed by Ardern *et al*¹ for the evaluation of return to sport outcomes after primary ACL reconstruction. The six items assessed were: selection criteria described, representative population selected, participants' pre-injury sports participation level reported, data collected prospectively, demographic data reported and postoperative sports participation level compared with pre-injury level. Items were scored as follows: one point for satisfaction of the criterion, no points for not satisfying the criterion. A score of zero was given if it was unclear whether the criterion had been satisfied or not. All articles were assessed independently by two assessors, disagreements being resolved by consensus, and third reviewer intervention in case of disagreement not being resolved. κ Coefficient for inter-rater agreement was calculated as well. The purpose of the quality assessment in this meta-analysis was to identify characteristics of study quality and design common to all included studies, and to provide a qualitative description of these characteristics in the meta-analysis.

Statistical analysis

Statistical evaluation was performed using MedCalc (MedCalc software, Acaciaaan 22, Ostend, Belgium). Data regarding return to sport, return to pre-injury sport level and return to high-level/competitive sports were combined via a proportion meta-analysis using a random effects model. I^2 index was used to measure the amount of inconsistency in the results of the included studies.^{19 20} A larger I^2 index indicates that a greater amount of the variability in the results is due to the heterogeneity rather than to chance.^{19 20} The mean and 95%CI was calculated for return to sport, return to pre-injury sport level and return to high level/competitive sport rates. Studies included in the meta-analyses were weighted according to sample size. Return to sport rate data from the individual studies were presented through forest-plot graphics.


The objective IKDC, KT-1000 or KT-2000 values and pivot-shift results were presented as the number of patients and percentage of patients for each category. The anteroposterior knee laxity measured with KT-1000 or KT-2000 was presented also as mean value expressed in mm.

RESULTS

The literature search yielded a total of 803 results. A flow diagram of the selection process is shown in [figure 1](#). After duplicates were removed, 472 studies remained. Of these, 414 were excluded after reviewing title and abstract review, as involving mostly surgical technique descriptions, in-vitro, biomechanics or radiologic studies, or because were studies not related to revision ACL reconstruction. The full text of the remaining 58 studies were obtained and reviewed for eligibility. Thirty-five studies were excluded because they did not report the outcomes requested in the inclusion criteria (24 studies) or because they were narrative or systematic reviews (11 papers). Therefore, 23 studies were included in the final analysis.^{10–13 21–39} A summary of the population examined with clinical outcomes is presented in [table 1](#).

Demographic data

The 23 reviewed studies presented the results of 1090 patients at a mean follow-up of 5.3 years (range 1.0–13.2 years). When

Figure 1 PRISMA flow diagram of the selection process.

gender was reported,^{10 11 13 24 26 28 29 33 37 38} revision ACL reconstruction was performed in 318 males (63%) and 187 females (27%). The mean age at revision surgery was 27.7 years (range 13–60 years), while the mean time from primary reconstruction and revision surgery, reported in 13 studies,^{10–12 22–24 27 28 30 31 33 34 37} was 43.8 months (range 1–276 months). Most of the studies reported the results of a first revision procedure, while only four studies^{12 29 34 38} reported at least one second-revision/third-revision procedure. Concerning the primary ACL lesions, 8% sustained ACL repair and 3% underwent an extra-articular procedure; when ACL reconstruction was instead performed, the primary graft to be used was bone patellar-tendon bone (BPTB) autograft in 57%, hamstrings autograft in 16%, other autografts in 2%, synthetic grafts in 7%, BPTB allograft in 2% and other allografts in 7%.

Regarding the revision procedure, eight studies reported the results of one single specific graft, another eight used a combination of different autografts (6 studies) or allografts (2 studies), while the remaining seven reported the use of a heterogeneous combination of allografts and autografts. Specifically, a BPTB autograft was used in 554 patients (50%), a hamstrings autograft in 326 (29%), a quadriceps tendon autograft in 71 (6%), a BPTB allograft in 98 (9%), an Achilles tendon allograft in 28 (3%) and other allografts in 38 (3%).

Quality assessment

Sixteen studies (69%) were prospective case series, 4 (17%) were retrospective case series while the remaining 3 (14%) were comparative studies between primary and revision reconstruction or between different populations.

Regarding quality assessment, only 5 studies (22%) fulfilled all the 6 items, while 9 (39%) fulfilled 5 items. Three studies (13%) fulfilled 4 items and 6 (26%) fulfilled 3 items (table 2).

The item “selection criteria described” was fulfilled by all 23 studies (100%), such as the “demographic data reported item”. The “representative population selected” item was fulfilled by six studies (26%) and the “data collected prospectively” item was fulfilled by 19 studies (83%). Finally, the item “participants’ pre-injury sports participation level reported” was fulfilled by 19 studies (83%), while the “postoperative sports participation level compared with pre-injury level” was fulfilled by 15 studies (65%).

The inter-rater agreement between the two reviewers regarding quality assessment was $k=0.92$ (95% CI 0.88 to 0.94).

The definitions of the parameter used to assess the return to sport and pre-injury sport level were not consistent between all the studies and are described in table 2.

Only six studies reported the outcomes of athletic populations, with inconsistent description of the practised sports; the other studies involved mixed populations of both athletic and non-athletic patients, or did not provide information regarding the populations studied.

Return to sport outcomes

Overall, 16 studies^{12 21–25 27–29 31 32 34–38} involving 543 patients reported a rate of return to sport activity at any level at the final follow-up; the pooled rate was 84% (CI 78 to 89) (figure 2), $I^2=65%$ ($p=0.0001$).

Fifteen studies,^{10 11 13 24 26–28 32–39} involving 790 patients, reported a rate of return to the same pre-injury sport level; the pooled rate was 52% (CI 36 to 67) (figure 3), $I^2=95%$ ($p=0.0001$).

Four studies,^{26 28 30 32} involving 186 patients, reported a rate of return to high-level/competitive sport; the pooled rate was 51% (CI 44 to 58) (figure 4), $I^2=0%$ ($p=0.3921$).

Only one study,¹² involving 11 multiple revisions performed with reharvested ipsilateral or contralateral BPTB, reported

Table 1 Details of patient populations, follow-up duration, time from primary to revision ACL reconstruction, objective outcomes at final follow-up (objective IKDC, difference of laxity between indexed and healthy knee and pivot-shift test) and return to sport outcomes for each study

Authors and year	Patients	Follow-up duration	Age at revision ACL reconstruction (ACLR)	Time from primary to revision ACL reconstruction (ACLR)	Final IKDC	Final laxity	Final pivot-shift n (%)	Return to sport n (%)
Noyes and Barber-Westin (1996) ²¹	65	3.5 years	25 years	NA	NA	3.4±4.8 mm	34 (52%) negative	51 (79%) Return to sport
						30 (53%) <3 mm 17 (30%) 3–5.5 mm 10 (17%) >5.5 mm	24 (37%) grade I 4 (6%) grade II 3 (5%) grade III	
Johnson <i>et al</i> (1996) ²²	22	2.3 years	27 years	NA	NA	1.9±4.4 mm	12 (60%) negative	16 (80%) Return to sport
						12 (66%) <3 mm 3 (17%) 3–5.5 mm 3 (17%) >5.5 mm	7 (35%) grade I 1 (5%) grade II 0 (0%) grade III	
Noyes and Barber-Westin (2001) ²³	25	2.3 years	25 years	30 months	NA	3.7±3.5 mm	20 (80%) negative— grade I	22 (88%) Return to sport
	22 M/3F					5 (20%) <2.5 mm 11 (44%) 3–5 mm 9 (36%) 5.5–10 mm	5 (20%) grade II 0 (0%) grade III	
O'Shea and Shelbourne (2002) ¹²	52 knees 51 patients	2.8 years	27 years	NA	NA	2.2±4.9 mm	NA	40 (72%) Return to sport
Taggart <i>et al</i> (2004) ²⁴	11	4.1 years	24 years	71 months	NA	1.6±1.2 mm	NA	11 (100%) Return to sport
						11 (100%) <3 mm		5.1±2.2 weeks to light sports 5.4±1.5 months to full sport
Carson <i>et al</i> (2004) ²⁵	20	3.4 years	30 years	49 months	NA	3 (15%) <1 mm 5 (25%) 1–3 mm 5 (25%) 3–5 mm 7 (35%) >5 mm	3 (15%) negative 9 (45%) grade I 4 (20%) grade II 4 (20%) grade III	8 (40%) Same pre-injury sport level 6 (30%) Lower level
	14 M/6F							
O'Neill (2004) ²⁶	43	Minimum 2 years	NA	NA	NA	2.9 mm 20 (47%) <3 mm 13 (30%) 3–4 mm 6 (14%) 4–5 mm 4 (9%) >5 mm	37 (86%) negative	12 (28%) Sport without limitation 20 (46%) Sport with limitation
	48 28 M/20F	7.5 years	33 years	60 months	20 (42%) A 20 (42%) B 6 (12%) C 2 (4%) D	10 (21%) 0 mm 22 (46%) 0–2 mm 3 (6%) 2–3 mm 10 (21%) 3–5 mm 3 (6%) >5 mm	NA	36 (75%) Same pre-injury sport level 21 (44%) High impact activities
Grossman <i>et al</i> (2005) ¹⁰	29	5.6 years	30.2 years	56 months	17 (58%) A 8 (28%) B 4 (14%) C 0 (0%) D	2.8 mm	27 (93%) negative 2 (7%) grade I 0 (0%) grade II 0 (0%) grade III	20 (68%) Same pre-injury sport level
	22 M/7F							

Continued

Table 1 Continued

Authors and year	Patients	Follow-up duration	Age at revision ACL reconstruction (ACLR)	Time from primary to revision ACL reconstruction (ACLR)	Final IKDC	Final laxity	Final pivot-shift n (%)	Return to sport n (%)
Thomas <i>et al</i> (2005) ¹³	49 37 M/12F	6.2 years	35.4 years	NA	12 (24%) A 28 (58%) B 8 (16%) C 1 (2%) D	1.9±2.4 mm 46 (94%) 0–2 mm 1 (2%) 3–4 mm 2 (4%) >5 mm	43 (88%) negative 5 (10%) grade I 1 (2%) grade II 0 (0%) grade III	0 (0%) Same pre-injury sport level
Ferretti <i>et al</i> (2006) ²⁷	28	5.0 years	34 years	60 months	15 (53%) A 11 (39%) B 2 (8%) C 0 (0%) D	2.5±1.8 mm 20 (71%) 0–3 mm 6 (21%) 3–5 mm 2 (8%) >5 mm	15 (54%) negative 11 (39%) grade I 2 (7%) grade II 0 (0%) grade III	6 (20%) Same pre-injury sport level 9 (30%) Lower level 30 (100%) Return to sport
Garofalo <i>et al</i> (2006) ²⁸	28 16 M/12F	4.2 years	27 years	26 months	5 (18%) A 21 (75%) B 2 (7%) C 0 (0%) D	3.1±2.0 mm 18 (65%) <3 mm 9 (32%) 3–5 mm 1 (3%) 6–10 mm 0 (0%) >10 mm	21 (75%) negative 7 (25%) grade I 0 (0%) grade II 0 (0%) grade III	26 (93%) Same pre-injury sport level 18 (64%) High competitive 6 (21%) Recreational 2 (7%) Light sport
Noyes and Barber-Westin (2006) ²⁹	21 15 M/6F	4.1 years	33 years	NA	5 (24%) A 11 (52%) B 3 (14%) C 2 (10%) D	2.1±2.2 mm 8 (38%) <3 mm 7 (33%) 3–5 mm 4 (19%) >5 mm	10 (48%) negative 7 (33%) grade I 3 (14%) grade II 1 (5%) grade III	15 (71%) Return to sport
Salmon <i>et al</i> (2006) ³⁰	50	7.4 years	27 years	36 months	33 (73%) A/B 12 (27%) C/D	2.5 mm 17 (50%) <3 mm 17 (50%) 3–5 mm 0 (0%) >5 mm	31 (69%) negative 14 (31%) grade I 0 (0%) grade II 0 (0%) grade III	26 (58%) Return to level I-II activity
Battaglia <i>et al</i> (2007) ³¹	63 36 M/27F	6.1 years	31 years	61 months	NA	3.9±2.4 mm 30 (48%) <3 mm 20 (31%) 3–5 mm 10 (21%) >5 mm	NA	42 (66%) Return to sport
Denti <i>et al</i> (2008) ³²	60	3.5 years	31 years	NA	18 (36%) A 23 (46%) B 9 (18%) C 0 (0%) D	28 (56%) <3 mm 17 (34%) 3–5 mm 5 (10%) 6–10 mm 0 (0%) >10 mm	NA	47 (78%) Same pre-injury sport level 8 (14%) Lower level 30 (50%) High competitive activities
Diamantopoulos <i>et al</i> (2008) ³³	107 64 M/43F	6.0 years	39 years	60 months	79 (74%) A 17 (16%) B 11 (10%) C 0 (0%) D	0.9±1.1 mm (KT)	79 (74%) negative 17 (16%) grade I 11 (10%) grade II 0 (0%) grade III	39 (36%) Same pre-injury sport level
Wegrzyn <i>et al</i> (2009) ³⁴	10 8 M/2F	3.2 years	30 years	26 months	4 (40%) A 5 (50%) B 1 (10%) C 0 (0%) D	1.3±1.9 mm 9 (90%) <5 mm 1 (10%) >5 mm	NA	1st revision: 5 (50%) Same pre-injury sport level 4 (40%) Lower level 2nd revision: 2 (20%) Same pre-injury sport level 4 (40%) Lower level

Continued

Table 1 Continued

Authors and year	Patients	Follow-up duration	Age at revision ACL reconstruction (ACLR)	Time from primary to revision ACL reconstruction (ACLR)	Final IKDC	Final laxity	Final pivot-shift n (%)	Return to sport n (%)
Muneta <i>et al</i> (2010) ³⁵	21 14 M/7F	3.3 years	27 years	NA	NA	1.7±1.8 mm	14 (78%) negative 3 (17%) grade I 1 (5%) grade II 0 (0%) grade III	13 (76%) Same pre-injury sport level 18 (86%) Return to sport
Reinhardt <i>et al</i> (2012) ³⁶	21	3.0 years	16.9 years	NA	NA	NA	20 (95%) negative 1 (5%) grade I 0 (0%) grade II 0 (0%) grade III	11 (52%) Same pre-injury sport level 20 (95%) Return to sport
Franceschi <i>et al</i> (2013) ³⁷	30 19 M/11F	6.8 years	29.1 years	34 months	27 (90%) A/B 3 (10%) C 0 (0%) D	3.1 mm	25 (83%) Negative 5 (17%) grade I 0 (0%) grade II 0 (0%) grade III	20 (67%) Same pre-injury sport level 7 (25%) Lower activity
Gifstad <i>et al</i> (2013) ¹¹	56 25 M/31F	7.5 years	26.5 years	38 months	NA	3.3±2.7 mm 28 (52%) <3 mm 17 (23%) 3–5 mm 8 (15%) >5 mm	45 (80%) negative— grade I 11 (20%) grade II-III	7 (13%) Same pre-injury sport level
Buda <i>et al</i> (2013) ³⁸	24 24 M/0F	3.3 years	NA	NA	4 (17%) A 16 (67%) B 2 (8%) C 2 (8%) D	3.1±1.1 mm	4 (17%) negative 18 (75%) grade I 2 (8%) grade II 0 (0%) grade III	17 (71%) Same pre-injury sport level 3 (13%) Lower level
Shelbourne <i>et al</i> (2014) ³⁹	259	7.2 years	22.4 years	NA	NA	2.3±1.8 mm	NA	178 (69%) Same pre-injury sport level

Pre-injury level is referred to the initial injury before primary ACL reconstruction. Values are expressed as the mean or absolute number and percentage, according to variable. F, females; M, males; IKDC, International Knee Documentation Committee; NA, not applicable.

Table 2 Details of study design, quality assessment, mixed or athletic population included in each study and the definition of return to sport extracted from the text of each study, with the scale to measure activity level (when used)

Authors and year	Study design	Quality assessment	Population	Return to sport evaluation
Noyes and Barber-Westin (1996) ²¹	Prospective observational study	5	Mixed	Sport activity preoperatively and at follow-up
Johnson <i>et al</i> (1996) ²²	Prospective observational study	5	Mixed	Sport activity frequency preoperatively and postoperatively
Noyes and Barber-Westin (2001) ²³	Prospective observational study	3	Mixed	Return to sports Cincinnati knee rating system
O'Shea and Shelbourne (2002) ¹²	Prospective observational study	3	Mixed	Return to sports
Taggart <i>et al</i> (2004) ²⁴	Retrospective case series	5	Athletes	Return to same pre-injury sport Return to sport at any level
Carson <i>et al</i> (2004) ²⁵	Retrospective case series	3	Mixed	Return to sports
O'Neill (2004) ²⁶	Prospective observational study	5	Mixed	Return to same pre-injury sport Return to pivoting/cutting/contact
Grossman <i>et al</i> (2005) ¹⁰	Retrospective case series	3	Mixed	Return to same pre-injury activity and sport Return to collegiate sport
Thomas <i>et al</i> (2005) ¹³	Prospective comparative study	4	Mixed	Return to original level of sport activity
Ferretti <i>et al</i> (2006) ²⁷	Prospective observational study	6	Athletes	Return to same pre-injury sports Return to sports
Garofalo <i>et al</i> (2006) ²⁸	Prospective observational study	6	Athletes	Return to previous sport activity IKDC activity score
Noyes and Barber-Westin (2006) ²⁹	Prospective observational study	3	Mixed	Return to sport Cincinnati knee rating system
Salmon <i>et al</i> (2006) ³⁰	Prospective observational study	4	Mixed	IKDC activity score
Battaglia <i>et al</i> (2007) ³¹	Prospective observational study	3	Mixed	Return to sport IKDC activity score
Denti <i>et al</i> (2008) ³²	Prospective observational study	5	Mixed	Return to same pre-injury sports
Diamantopoulos <i>et al</i> (2008) ³³	Prospective observational study	5	Mixed	Return to same pre-injury sport activity
Wegrzyn <i>et al</i> (2009) ³⁴	Retrospective case series	4	Mixed	Return to same pre-injury sports
Muneta <i>et al</i> (2010) ³⁵	Retrospective comparative study	5	Mixed	Return to sport activity equal to pre-injury level
Reinhardt <i>et al</i> (2012) ³⁶	Prospective observational study	6	Athletes	Return to same or higher activity/sport level IKDC activity score
Franceschi <i>et al</i> (2013) ³⁷	Prospective observational study	5	Mixed	Return to pre-injury sport level Return to lower non-impact sports
Gifstad <i>et al</i> (2013) ¹¹	Prospective observational study	5	Mixed	Return to same or better pre-injury sport level
Buda <i>et al</i> (2013) ³⁸	Prospective observational study	6	Athletes	Return to same pre-injury sport level Sport resumption at lower level
Shelbourne <i>et al</i> (2014) ³⁹	Prospective comparative study	6	Athletes	Return to same sport at previous level

IKDC, International Knee Documentation Committee.

the time to return to sport. Return to light sport was achieved after 5.1 ± 2.2 weeks after revision, while return to full sport was achieved after 5.4 ± 1.5 months.

Only three studies,^{21 28 31} involving 174 patients, reported the main reason for not returning to sport after ACL revision surgery: 31 patients (69%) were not able to return to sport because of knee-related problems, 10 (22%) because of the fear of re-injury while 4 (9%) were unable to do so for other reasons.

Secondary outcomes

Twelve studies^{10 13 26–30 32–34 37 38} reported the IKDC results in 392 patients. At final follow-up, 177 patients (45%) had knee function rated as normal (A), 160 (41%) as nearly normal (B), 48 (12%) as abnormal (C) and 7 (2%) as severely abnormal (D).

Twenty-two studies^{10–13 21–35 37–39} reported laxity evaluation at the final follow-up of 1069 patients, reporting a mean final value of 2.6 mm (range -4 –6 mm). When reported, 325 patients (58%) presented a value of side-to-side difference at manual maximum displacement test < 3 mm, 169 (30%) between 3 and 5 mm, and 69 (12%) > 5 mm.

Sixteen studies^{10 11 13 21 22 24 26–30 33 35–38} reported the results of the postoperative Pivot-Shift test in 515 patients. The

test was considered negative in 338 patients (67%), while it was graded as grade I in 140 (26%), grade II in 29 (5%) and grade III in 8 (2%).


Therefore, 337/392 (86%) patients presented a normal or nearly normal knee (group A and B according to objective IKDC), 494/563 (88%) had normal values of static anteroposterior laxity (side to side difference < 5 mm at KT-1000/2000) and 478/515 (93%) had a normal or nearly normal rotatory laxity (normal or grade I pivot-shift test).

DISCUSSION

The main findings of this meta-analysis are that 84% of patients returned to sport after revision ACL reconstruction, while 52% returned to the same pre-injury sport level and 51% returned to high-level/competitive sports.

The return to sport after both primary and revision ACL reconstruction represents a complex outcome to evaluate. First of all, a great inconsistency in reporting the results was found in most of the studies, as a clear and unequivocal definition of return to sport was not provided. For example, the concept of "return to pre-injury sport level" presented a not unequivocal interpretation in most of the studies, especially regarding the kind of sport and its level (amateur, competitive, elite), possibly

Figure 2 Forest-plot graphic of return to sport outcome; %, mean percentage; CI, 95% CIs.



contributing to the production of high variability of the results reported in this meta-analysis. In fact, the latter outcome was mostly assessed with the sentence “patients returned to the same pre-injury sport activity” or “were able to perform sport equal to pre-injury level”, without mentioning the type of sport, level and intensity. Moreover, only a few studies reported the specific sport performed by each of the patients, thus possibly creating a bias because it could be speculated that patients who perform less demanding activities like tennis or running could hypothetically return to sport easier than patients performing pivoting sports like soccer, basketball and handball.


With the available data obtained from this meta-analysis, the return to any sport outcome was similar to what was reported after primary ACL reconstruction;¹ while the return to pre-injury sport level resulted lower compared to primary ACL reconstruction. This could also be due to the higher age at revision ACL reconstruction. In fact, the revision ACL reconstruction could be performed in patients near to 40 years of age. Therefore it could be difficult for them to return to perform the same activity of long time before, when they had the primary ACL injury, and this is not always necessarily due to reasons related to knee performance. Moreover it should be noted that the activity before a primary ACL lesion was used as a landmark in most of the studies. Conversely, the unreported activity level prior to primary ACL reconstruction failure would have probably pointed out the real impact of revision procedure on sport

activity, or at least provided adjunctive information regarding the changes of sport practice along the whole follow-up from the very first injury to final evaluation.

Regarding the impact of graft type in the return to sport outcomes, the analysis of the data from the current literature produced inconsistent results. This is mostly due to the lack of well-designed RCTs and to the high heterogeneity in graft choice, between and within each single study. This situation reflects the complexity of revision surgery, such as graft availability derived from previous reconstructions, the status of the contralateral knee, the availability of allograft tissue or the need for a graft with bone stock, which often make the graft choice a necessity rather than a preference. Most of authors used the unharvested BPTB or hamstrings as first choice, while reserving quadriceps tendon autograft, contralateral grafts or various allografts only in case of repeated revision, combined posterior cruciate ligament or posterolateral corner reconstruction, previous multiligament reconstruction or other particular extraordinary circumstances. Therefore only a few studies included revision ACL reconstruction with a single specific graft, making it impossible to compare sport-specific outcomes based on graft choice.

A wide discrepancy is noted between successful results of revision ACL reconstruction based on objective evaluation and sport-related outcomes; normal or nearly normal results were reported in 86%, 88% and 93% for objective IKDC, anteroposterior knee laxity and Pivot-Shift test, respectively, against a rate

Figure 3 Forest-plot graphic of return to pre-injury sport level; %, mean percentage; CI, 95% CIs.






Figure 4 Forest-plot graphic of return to high-level/competitive sport; %, mean percentage; CI, 95% CIs.

of 52% of return to pre-injury sport level. The measurement of sport participation represents a relevant issue, especially in athletes and high-level patients, probably superior to objective tools such as knee stability, ROM and muscle strength, because sport participation is more meaningful for them in their daily lives. The reasons for the discrepancy between objective and sport-related outcomes should be evaluated and addressed, as probably in not all cases the inability to return to the same pre-injury sport level was due to the indexed knee problems. In contrast, fear of re-injury, priorities and personality could influence the decision to return to sport.^{3 40} It has in fact been previously demonstrated that issues unrelated to surgical outcomes, such as lack of motivations, reduced confidence, fear of sport incompetence or lower priority on sport due to family/work-related reasons, have negative effects on the return to pre-injury sport level.³ Such evidence, besides representing another potential contributing factor to the high variability of results, could point out the necessity of psychological interventions integrated in the postoperative rehabilitation process to possibly try to improve the outcomes of revision ACL reconstruction, as already suggested for primary ACL reconstruction.⁴¹ Moreover, poor surgery, inadequate rehabilitation and impairment of muscle performance, isokinetic strength and functional tests could also affect the ability to return to sport.^{4 5}

This meta-analysis presents some strong limitations. First of all, it ought to be mentioned that the return to sport rate considered in this study actually represents the number of patients who are still able to perform sports at the final follow-up. It is plausible that, during the considered follow-up, some patients were able to return to sport after revision ACL reconstruction and rehabilitation, but that several reasons caused them to eventually abandon the practice shortly afterwards. The actual return to sports rate would thus include all patients/athletes being able to resume sports at any time after ACL revision surgery, irrespective of whether they were still active or not at the time of follow-up. However, the 'return to sports rate' and the 'still active at follow-up rate' were not reported separately in the included studies, thus making an adequate analysis impossible as well as potentially generating misinterpretation of the results. Therefore, the reader should be aware of this major issue and take a conservative approach when drawing conclusions from this meta-analysis.

Another major limitation was the high heterogeneity of the studies included, particularly regarding patients' characteristics and surgical treatment, which did not allow an analysis of subgroups regarding sport participation and objective outcomes. This is true also for first or second or third revision ACL reconstruction, as only two studies exclusively evaluated this population, making further analysis impossible. Furthermore, the dishomogeneous and equivocal definition of return to sport and

return to pre-injury sport level could have created a bias when interpreting the results. Also, owing to the lack of studies with a follow-up of less than 2 years, it was impossible to compare the short-term results of revision ACL reconstruction with the medium-term and long-term results, as already performed by Ardern *et al*¹ when evaluating primary ACL reconstruction. This made it impossible to monitor the sport practice from short-term to long-term follow-up, and to detect potential decreases of sport activity with the passage of time. Finally, the evaluation of subjectively measured parameters, such as Pivot-shift test and IKDC score, could have produced a concrete bias regardless of whether independent examiners were involved in the patient's evaluation, because these objective measures could lack sensitivity to detect subtle differences in function after revision ACL reconstruction. Unfortunately, this issue could not have been investigated adequately due to the great inconsistency in reporting the details regarding the examiner(s) in each study. On the other hand, the evaluation of an objective parameter such as knee laxity determined with KT-1000/2000, despite the lack of inter-rater reliability across different studies, could provide a concrete measure of postoperative outcomes.

Future high-quality RCTs or prospective case series with homogeneous populations, especially regarding graft choice and surgical techniques, could improve the knowledge and understanding of revision ACL reconstruction outcomes. Moreover, the use of a clear definition of sport-related outcomes, such as return to pre-injury sport level, return to training or competitive sport, explicit presentation of the type of sport and level practised before ACL injury, at the end of rehabilitation, and at the final follow-up, could avoid a misleading interpretation of revision ACL reconstruction outcomes and better understanding of the effective ability to perform sport activity after this kind of surgical procedure.

What is known about the topic?

- ▶ Revision ACL reconstruction is an increasingly performed procedure due to the high number of primary ACL reconstructions and failures.
- ▶ Revision ACL reconstruction is able to restore knee stability and improve knee function.
- ▶ Return to sport after revision ACL reconstruction has not been deeply investigated.
- ▶ Wide ranges of return to sport are present in the current literature.

What are the new findings?

- ▶ 84% of patients who had revision ACL reconstruction returned to any kind of sport activity.
- ▶ 52% of patients returned to their pre-injury sport level.
- ▶ There was a higher percentage of patients with normal or nearly normal knees at final follow-up, compared to the proportion who returned to sport.
- ▶ A homogeneous definition of return to sport when dealing with revision ACL reconstruction needs to be clarified in future studies.

CONCLUSIONS

On the basis of the data of the present meta-analysis, 84% of patients returned to sport after revision ACL reconstruction, while 52% returned to the same pre-injury sport level. Normal, or nearly normal, objective results were reported in 86%, 88% and 93% for objective IKDC, anteroposterior knee laxity and Pivot-Shift test, respectively. The high heterogeneity of patient populations and sport-related outcome definitions represent the major biases when evaluating return to sport outcomes after ACL reconstruction outcomes.

Contributors AG, GMMM and SZ performed a database search and review of articles. MPN and SDV performed data collection and an analysis of the results. MM wrote the paper and critically evaluated the results.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Ardern CL, Webster KE, Taylor NF, *et al.* Return to sport following anterior cruciate ligament reconstruction surgery: a systematic review and meta-analysis of the state of play. *Br J Sports Med* 2011;45:596–606.
- Ardern CL, Taylor NF, Feller JA, *et al.* Psychological responses matter in returning to preinjury level of sport after anterior cruciate ligament reconstruction surgery. *Am J Sports Med* 2013;41:1549–58.
- Tjong VK, Murnaghan ML, Nyhof-Young JM, *et al.* A qualitative investigation of the decision to return to sport after anterior cruciate ligament reconstruction: to play or not to play. *Am J Sports Med* 2014;42:336–42.
- Thomé R, Kaplan Y, Kvist J, *et al.* Muscle strength and hop performance criteria prior to return to sports after ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2011;19:1798–805.
- Bizzini M, Silvers HJ. Return to competitive football after major knee surgery: more questions than answers? *J Sports Sci* 2014;32:1209–16.
- Kvist J. Rehabilitation following anterior cruciate ligament injury: current recommendations for sports participation. *Sports Med* 2004;34:269–80.
- Warner SJ, Smith MV, Wright RW, *et al.* Sport-specific outcomes after anterior cruciate ligament reconstruction. *Arthroscopy* 2011;27:1129–34.
- Wright RW, Gill CS, Chen L, *et al.* Outcome of revision anterior cruciate ligament reconstruction: a systematic review. *J Bone Joint Surg Am* 2012;94:531–6.
- Gianotti SM, Marshall SW, Hume PA, *et al.* Incidence of anterior cruciate ligament injury and other knee ligament injuries: a national population-based study. *J Sci Med Sport* 2009;12:622–7.
- Grossman MG, ElAttrache NS, Shields CL, *et al.* Revision anterior cruciate ligament reconstruction: three- to nine-year follow-up. *Arthroscopy* 2005;21:418–23.
- Gifstad T, Drogset JO, Viset A, *et al.* Inferior results after revision ACL reconstructions: a comparison with primary ACL reconstructions. *Knee Surg Sports Traumatol Arthrosc* 2013;21:2011–18.
- O'Shea JJ, Shelbourne KD. Anterior cruciate ligament reconstruction with a reharvested bone-patellar tendon-bone graft. *Am J Sports Med* 2002;30:208–13.
- Thomas NP, Kankate R, Wandless F, *et al.* Revision anterior cruciate ligament reconstruction using a 2-stage technique with bone grafting of the tibial tunnel. *Am J Sports Med* 2005;33:1701–9.
- Moher D, Liberati A, Tetzlaff J, *et al.*, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- Hefti F, Müller W, Jakob RP, *et al.* Evaluation of knee ligament injuries with the IKDC form. *Knee Surg Sports Traumatol Arthrosc* 1993;1:226–34.
- Wera JC, Nyland J, Ghazi C, *et al.* International knee documentation committee knee survey use after anterior cruciate ligament reconstruction: a 2005–2012 systematic review and world region comparison. *Arthroscopy* 2014;30:1505–12.
- Jakob RP, Stäubli HU, Deland JT. Grading the pivot shift. Objective tests with implications for treatment. *J Bone Joint Surg Br* 1987;69:294–9.
- Musahl V, Hoshino Y, Ahlden M, *et al.* The pivot shift: a global user guide. *Knee Surg Sports Traumatol Arthrosc* 2012;20:724–31.
- Hedges LV, Vevea JL. Fixed-and random-effects models in meta-analysis. *Psychol Methods* 1998;3:486–504.
- Higgins JP, Thompson SG, Deeks JJ, *et al.* Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- Noyes FR, Barber-Westin SD. Revision anterior cruciate ligament surgery: experience from Cincinnati. *Clin Orthop Relat Res* 1996;325:116–29.
- Johnson DL, Swenson TM, Irrgang JJ, *et al.* Revision anterior cruciate ligament surgery: experience from Pittsburgh. *Clin Orthop Relat Res* 1996;325:100–9.
- Noyes FR, Barber-Westin SD. Revision anterior cruciate surgery with use of bone-patellar tendon-bone autogenous grafts. *J Bone Joint Surg Am* 2001;83:1131–43.
- Taggart TF, Kumar A, Bickerstaff DR. Revision anterior cruciate ligament reconstruction: a midterm patient assessment. *Knee* 2004;11:29–36.
- Carson EW, Anisko EM, Restrepo C, *et al.* Revision anterior cruciate ligament reconstruction: etiology of failures and clinical results. *J Knee Surg* 2004;17:127–32.
- O'Neill DB. Revision arthroscopically assisted anterior cruciate ligament reconstruction with previously unharvested ipsilateral autografts. *Am J Sports Med* 2004;32:1833–41.
- Ferretti A, Conteduca F, Monaco E, *et al.* Revision anterior cruciate ligament reconstruction with doubled semitendinosus and gracilis tendons and lateral extra-articular reconstruction. *J Bone Joint Surg Am* 2006;88:2373–9.
- Garofalo R, Djahangiri A, Siegrist O. Revision anterior cruciate ligament reconstruction with quadriceps tendon-patellar bone autograft. *Arthroscopy* 2006;22:205–14.
- Noyes FR, Barber-Westin SD. Anterior cruciate ligament revision reconstruction: results using a quadriceps tendon-patellar bone autograft. *Am J Sports Med* 2006;34:553–64.
- Salmon LJ, Pinczewski LA, Russell VJ, *et al.* Revision anterior cruciate ligament reconstruction with hamstring tendon autograft: 5- to 9-year follow-up. *Am J Sports Med* 2006;34:1604–14.
- Battaglia MJ, Ilnid, Cordasco FA, Hannafin JA, *et al.* Results of revision anterior cruciate ligament surgery. *Am J Sports Med* 2007;35:2057–66.
- Denti M, Lo Vetere D, Bait C, *et al.* Revision anterior cruciate ligament reconstruction: causes of failure, surgical technique, and clinical results. *Am J Sports Med* 2008;36:1896–902.
- Diamantopoulos AP, Lorbach O, Paessler HH. Anterior cruciate ligament revision reconstruction: results in 107 patients. *Am J Sports Med* 2008;36:851–60.
- Wegrzyn J, Chouteau J, Philippot R, *et al.* Repeat revision of anterior cruciate ligament reconstruction: a retrospective review of management and outcome of 10 patients with an average 3-year follow-up. *Am J Sports Med* 2009;37:776–85.
- Muneta T, Hara K, Ju YJ, *et al.* Revision anterior cruciate ligament reconstruction by double-bundle technique using multi-strand semitendinosus tendon. *Arthroscopy* 2010;26:769–81.
- Reinhardt KR, Hammoud S, Bowers AL, *et al.* Revision ACL reconstruction in skeletally mature athletes younger than 18years. *Clin Orthop Relat Res* 2012;470:835–42.
- Franceschi F, Papalia R, Del Buono A, *et al.* Two-stage procedure in anterior cruciate ligament revision surgery: a five-year follow-up prospective study. *Int Orthop* 2013;37:1369–74.
- Buda R, Ruffilli A, Di Caprio F, *et al.* Allograft salvage procedure in multiple-revision anterior cruciate ligament reconstruction. *Am J Sports Med* 2013;41:402–10.
- Shelbourne KD, Benner RW, Gray T. Return to sports and subsequent injury rates after revision anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med* 2014;42:1395–400.
- Kvist J, Ek A, Sporrstedt K, *et al.* Fear of re-injury: a hindrance for returning to sports after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2005;13:393–7.
- Zaffagnini S, Russo RL, Marcheggiani Muccioli GM, *et al.* The Videoinight® method: improving rehabilitation following anterior cruciate ligament reconstruction—a preliminary study. *Knee Surg Sports Traumatol Arthrosc* 2013;21:851–8.



After revision anterior cruciate ligament reconstruction, who returns to sport? A systematic review and meta-analysis

Alberto Grassi, Stefano Zaffagnini, Giulio Maria Marcheggiani Muccioli, Maria Pia Neri, Stefano Della Villa and Maurilio Marcacci

Br J Sports Med published online June 10, 2015

Updated information and services can be found at:

<http://bjsm.bmj.com/content/early/2015/06/10/bjsports-2014-094089>

These include:

References

This article cites 41 articles, 18 of which you can access for free at: <http://bjsm.bmj.com/content/early/2015/06/10/bjsports-2014-094089#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[BJSM Reviews with MCQs](#) (88)

[Knee laxity](#) (7)

[Procedures](#) (44)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>